



LYMPHA PRESS

ORDER FORM FAX #: 800.758.0339

QUESTIONS CALL: 800.734.0422

Proud to offer Lympha Press and to Serve You and Your Patients with Communication, Care, and Quality

DATE OF ORDER \_\_\_/\_\_\_/\_\_\_

Patient Name: \_\_\_\_\_ DOB \_\_\_/\_\_\_/\_\_\_

Patient Phone Number: \_\_\_\_\_

Extremities for Treatment: LEGS: \_\_\_LT \_\_\_RT \_\_\_ With Abdomen (Lympha Pants) ARMS: \_\_\_LT \_\_\_RT \_\_\_ With Trunk (Comfy Sleeve)

Special request: \_\_\_\_\_

TREATMENT: Pressures \_\_\_\_\_ Frequency \_\_\_\_\_/DAY \_\_\_\_\_MIN Length of Necessity: \_\_\_\_\_months (99 = lifetime)

Default Settings: 40mmHg, TID/BID, 60 min

DIAGNOSIS:

- [ ] Q82.0 HEREDITARY LYMPHEDEMA
[ ] I89.0 SECONDARY LYMPHEDEMA DUE TO:
[ ] VENOUS INSUFFICIENCY
[ ] TUMOR(S)
[ ] SCARRING OF THE LYMPH CHANNELS FROM CELLULITIS AND/OR LYMPHANGITIS
[ ] CANCER SURGERY AND/OR RADIATION
[ ] OTHER
[ ] I97.2 POST MASTECTOMY SYNDROME
[ ] I87.2 CHRONIC VENOUS INSUFFICIENCY
[ ] Varicose vein with ulcer
[ ] Venous hypertension with an ulcer

PRESCRIBING PHYSICIAN: (please print)

LAST \_\_\_\_\_

FIRST \_\_\_\_\_

NPI#: \_\_\_\_\_

PHONE \_\_\_\_\_

FAX \_\_\_\_\_

E-MAIL \_\_\_\_\_

PHYSICIAN SIGNATURE \_\_\_\_\_

IN CLINIC CONTACT: \_\_\_\_\_

DATE \_\_\_/\_\_\_/\_\_\_



## Please Include with Order:

- **FAX COVER SHEET** with contact person name and phone/fax number
- **PATIENT DEMOGRAPHICS** including insurance information
- **LAST 4+ WEEKS OF CHART NOTES** (*if available*)  
(see documentation examples below)

## Documentation Examples:

### 1. DIAGNOSIS

**LYPHHEDEMA**, identified as one of the following:

- Hereditary (Q82.0)
- Secondary (I89.0)
- Post Mastectomy (I97.2)

**CHRONIC VENOUS INSUFFICIENCY** (I87.2)  
with **VENOUS STASIS ULCER(s)** (I83.0 or I87.31)

### 2. SEVERITY, INCLUDING SPECIFIC EXAMPLES:

**STAGE 1, 2, or 3 LYPHHEDEMA**

Symptom examples include:

Skin changes (dermal metaplasia)	Fibrosis
Pitting or non-pitting edema	Hyperkeratosis
Papillomatosis cutis lymphostatica	Hyperpigmentation
Skin breakdown with persisting fluid leakage	
Elephantiasis ...etc.	

### 3. CONSERVATIVE TREATMENTS (Spanning 4 Weeks)

1. Regular Elevation
2. Regular Exercise
3. Compression Wrapping or Class 2 Compression Sleeve/Stocking

### 4. SYMPTOMS PERSIST

## Sample Dictations

*"Patient presents with Stage 1, 2, or 3 lymphedema. Recommend regular elevation, exercise, and Class 2 compression stockings (or wrapping system applied)"*

- **After 4 weeks (28 days)** -

*"Despite regular and compliant use of conservative treatments, including exercise, elevation, and Class 2 compression stockings, the patient still presents with Stage 1, 2, or 3 lymphedema."*

## LYPHA PRESS CONTRAINDICATIONS:

Use of the Lympha Press® system is not recommended in the presence of one or more of the following conditions:

- Known or suspected deep vein thrombosis (DVT) or pulmonary embolism
- During the inflammatory phlebitis process
- Acute infection of the affected limb
- Decompensated cardiac failure
- Severe arteriosclerosis or other ischemic vascular disease
- Any circumstance where increased venous and lymphatic return is undesirable

Due to movement of fluids in the body when using the system, exercise caution when using on patients with heart disease. High pressure is not recommended for patients who have peripheral occlusion disease. The abdominal area should not be treated during pregnancy.