

Therapy Outcomes in Post-Acute Care Settings (TOPS) Study Chartbook

April 2021

Jointly Commissioned by



Table of Contents

Background & Objectives	3
How do Beneficiaries Compare Across Settings?	9
Key Finding #1	10
How Does Intensity of OT & PT Services Relate to Functional Status & Readmissions?	20
Key Finding #2	21
Key Finding #3	22
Setting-Specific Findings	
Home Health Agencies	23
Skilled Nursing Facilities	30
Inpatient Rehabilitation Facilities	37
Discussion & Policy Implications	44
Appendix	49

Background & Objectives

Background

Occupational therapy and physical therapy services are a critical component of post-acute care (PAC).

New and emerging payment policies have the potential to impact therapy provision in PAC settings, including:



**Home Health
Agencies**



**Skilled Nursing
Facilities**




**Inpatient Rehab
Facilities**

The study was conducted by Dobson DaVanzo and Associates, LLC in 2018-2020 using 2015 & 2016 Medicare claims and administrative data. The study included Medicare fee-for-service beneficiaries who received OT and/or PT services in the first PAC stay following a hospitalization.

Objective #1: Beneficiaries Served in PAC Settings

How do Beneficiaries Compare Across Settings?


Race & Ethnicity



Dual-Eligible Status




Gender



ICU Use & Comorbidities



Age



Diagnostic Groups



Objective #2: Service Intensity & Outcomes

How Does Intensity of OT & PT Services Relate to:

1 Change in functional status during first PAC episode?

Functional status includes ability to perform:

Dressing	Walking/ Locomotion	Toileting
Bathing	Transferring	Feeding

A score of 0 is dependent on all 6 items. A score of 36 is independent on all 6 items.

Objective #2: Service Intensity & Outcomes

How Does Intensity of OT & PT Services Relate to:

2 Readmission to acute care within 30 days from PAC discharge?

All models adjusted/controlled for:

Admitted to PAC 5 Days or Less After Hospital Discharge	PAC Interrupted by Acute Admission	Alzheimer's and/or Dementia	Depression
Hospital Length of Stay	Use of ICU in Hospital	Stineman Stage III ADL	Comorbid Conditions
Age	Gender	ESRD	Rural
Race/Ethnicity	Dual Beneficiary	Disability Status	State

Number of home health visits (for home health model only).

Study Population

1.4 million

Medicare PAC stays with OT and/or PT services following hospitalization in 2015 and 2016



552,000+
**Home Health
Stays**



693,000+
**Skilled Nursing
Stays**



197,000+
**Inpatient Rehab
Stays**

Key Findings:


How Do Beneficiaries Compare Across Settings?

Key Finding #1

Distinct people with **complex** conditions & comorbidities receive occupational therapy and physical therapy in each post-acute care setting.

Race & Ethnicity

Race & ethnicity of people receiving therapy services were mostly consistent in each setting.



Home Health




Skilled Nursing



Inpatient Rehab


Gender

In all settings, the majority of people receiving therapy was female.



Age Groups by Setting

The largest age group receiving therapy in HH agencies was between 65 and 74 years old, in IRFs was between 75 and 84 years old, and in SNFs was 85 or older.



Home Health




Skilled Nursing



Inpatient Rehab

Medicare & Medicaid Dual Eligible

SNFs had a larger proportion than did HH and IRFs of dual-eligible people receiving therapy services.



Home Health




Skilled Nursing



Inpatient Rehab

Alzheimer's Disease and/or Dementia

Nearly half of people in SNFs who receive therapy have dementia, compared with 20% in HH and 31% in IRFs.



Home Health




Skilled Nursing



Inpatient Rehab

Depression

Depression was common in all settings. More than 40% of people receiving therapy in SNFs and IRFs had a diagnosis of depression.



Home Health




Skilled Nursing



Inpatient Rehab

ICU Use During Hospitalization

Nearly 1 in 4 people receiving therapy in HH were admitted to the ICU during hospitalization. Nearly 30% of those in SNFs and 40% in IRF spent time in the ICU.



Home Health




Skilled Nursing



Inpatient Rehab

Major Diagnoses Across Settings

Although all major diagnostic groups are served in each setting, some groups are more prevalent in specific settings. Nearly half of people with stroke receive care in an IRF. Nearly half of people with congestive heart failure receive care in a SNF. Almost two-thirds of people with joint replacement received HH care.



Home Health




Skilled Nursing



Inpatient Rehab

Functional Status at Admission

The median functional status of people admitted to PAC is highest (meaning most independent) in HH and lowest in IRFs.



Home Health



Skilled Nursing



Inpatient Rehab

Key Findings:

How Does Intensity of OT & PT Services Relate to Functional Status & Readmissions?

Key Finding #2

The intensity of occupational therapy and physical therapy services is associated with **improved ability** to perform **everyday functional activities** such as bathing, dressing, and transferring.

Key Finding #3

People who received the **fewest minutes** of physical therapy and occupational therapy **were at the highest risk** for hospital **readmission** across all post-acute care settings.

These findings were consistent for patients with medical and surgical conditions, suggesting a robust impact across Medicare populations.

Setting-Specific Findings

Home Health Agencies



Home Health Overview


Group	Total Hours of OT & PT			
	All Diagnoses	Stroke	CHF	Joint Replacement
Low-Intensity Group	Less than 1.6 hours	Less than 1.9 hours	Less than 1 hour	Less than 3.8 hours
Typical-Intensity Group	5.3 to 6 hours	6 to 7.3 hours	4.5 to 5.5 hours	6.3 to 7 hours
High-Intensity Group	12.5 to 30 hours	16.5 to 29.3 hours	12 to 21.3 hours	12.3 to 30 hours

People who received the **least** therapy across all HH stays are in the low-intensity group. People who received the **most** therapy across HH stays are in the high intensity group. Each group represents 10% of total stays. The low-intensity and high-intensity groups represent the low and high extremes of therapy.

Readmissions After Home Health: All Diagnoses

People in the low-intensity group had the highest readmission risk in HH.


30-Day Readmission Rate After HH Discharge: All Diagnoses



Change in Function in Home Health: All Diagnoses

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


**Function Status Change From Admission to Discharge:
All Diagnoses**



Change in Function in Home Health: Joint Replacement

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


**Function Status Change From Admission to Discharge:
Diagnosis of Joint Replacement**



Change in Function in Home Health: Congestive Heart Failure

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: Diagnosis of CHF



Change in Function in Home Health: Stroke

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.

Function Status Change from Admission to Discharge: Diagnosis of Stroke



Setting-Specific Findings

Skilled Nursing Facilities



Skilled Nursing Facilities Overview


Decile	Total Hours of OT & PT			
	All Diagnoses	Stroke	CHF	Joint Replacement
Low-Intensity Group	Less than 6.3 hours	Less than 9.3 hours	Less than 6.9 hours	Less than 3.7 hours
Typical-Intensity Group	22.3 to 27.6 hours	29.6 to 39 hours	21.9 to 26.5 hours	13.7 to 17.5 hours
High-Intensity Group	90.1 to 219 hours	128 to 219.3 hours	80.1 to 165 hours	58.3 to 151.9 hours

People who received the **least** therapy across all SNF stays are in the low-intensity group. People who received the **most** therapy across SNF stays are in the high-intensity group. Each group represents 10% of total stays. The low-intensity and high-intensity groups represent the low and high extremes of therapy.

Readmissions After Skilled Nursing Facility Stay: All Diagnoses

The high-intensity group had lowest readmission followed by typical intensity. The low-intensity group had the highest readmission rate.


30-Day Readmission Rate After SNF Discharge: All Diagnoses



Change in Function in Skilled Nursing Facility Stay: All Diagnoses

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: All Diagnoses



Change in Function in Skilled Nursing Facility Stay: Stroke

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: Diagnosis of Stroke



Change in Function in Skilled Nursing Facility Stay: Congestive Heart Failure

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: Diagnosis of CHF



Change in Function in Skilled Nursing Facility Stay: Joint Replacement

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.

Function Status Change From Admission to Discharge: Diagnosis of Joint Replacement



Setting-Specific Findings

Inpatient Rehabilitation Facilities



Inpatient Rehabilitation Facilities Overview


Group	Total Hours of OT & PT			
	All Diagnosis	Stroke	CHF	Joint Replacement
Low-Intensity Group	Less than 13 hours	Less than 15.2 hours	Less than 12.9 hours	Less than 12.4 hours
Typical-Intensity Group	21.8 to 24.4 hours	27 to 30.6 hours	20.5 to 22.5 hours	19.2 to 21.2 hours
High-Intensity Group	41.1 & ≤85	>54 & ≤85	>34.7 & ≤55.7	>33.3 & ≤50.2

People who received the **least** therapy across all IRF stays are in the low-intensity group. People who received the **most** therapy across IRF stays are in the high-intensity group. Each group represents 10% of total stays. The low-intensity and high-intensity groups represent the low and high extremes of therapy.

Readmissions After Inpatient Rehabilitation Facility Stay: All Diagnoses

People who received the least OT & PT had the highest likelihood of readmission in IRF.


30-Day Readmission Rate After IRF Discharge: All Diagnoses



Change in Function in Inpatient Rehabilitation Facility Stay: All Diagnoses

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: All Diagnoses



Change in Function in Inpatient Rehabilitation Facility Stay: Stroke

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: Diagnosis of Stroke



Change in Function in Inpatient Rehabilitation Facility Stay: Congestive Heart Failure

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.


Function Status Change From Admission to Discharge: Diagnosis of CHF



Change in Function in Inpatient Rehabilitation Facility Stay: Joint Replacement

Both typical and high-intensity groups had significantly more functional improvement than the low-intensity group.

Function Status Change From Admission to Discharge: Diagnosis of Joint Replacement



Discussion & Policy Implications

OT & PT Are Related to Improved Function

The study provides clear evidence that physical therapy and occupational therapy services improve patient outcomes across all PAC settings.

Findings consistently indicate that these services are associated with greater increases in function, suggesting a high potential for harm to patients who receive the fewest minutes of therapy.

These findings highlight the importance of matching delivery of therapy services to patient needs.

OT & PT Are Related to Reduced Readmission

Thirty-day readmission rates to acute care hospitals following a PAC episode decreased after therapy and appear to have varying threshold effects within each setting.

Readmission rates help to reflect whether patients are receiving appropriate levels of care following discharge from the hospital.

Patients who receive physical therapy and occupational therapy during their initial PAC episode are less likely to be readmitted and more likely to have better outcomes, such as survival, functional ability, quality of life, and participation in daily living activities.

Policy Implications

The results of this report illustrate that the U.S. Centers for Medicare & Medicaid Services (CMS) should approach the research and development of a unified PAC prospective payment system slowly and thoughtfully.

Findings from the TOPS study provide strong evidence of the distinctions among post-acute care settings, with different patterns of rehabilitation delivered to patients who vary in needs and complexities in each setting.

Policymakers should not assume that there is clear overlap among the PAC settings for treating similar patients, as patients in each setting have different treatment goals and are at different points in their recovery.

These findings also indicate that CMS should monitor the level of therapy provided in PAC settings as payment incentives change in order to help protect patient access to medically necessary skilled therapy, especially patients with high rehabilitation needs.

For More Information

[APTA: advocacy@apta.org](mailto:advocacy@apta.org)

[AOTA: regulatory@aota.org](mailto:regulatory@aota.org)

The American Physical Therapy Association and the American Occupational Therapy Association jointly commissioned Dobson DaVanzo and Associates to examine provision of Medicare occupational therapy and physical therapy services within inpatient rehabilitation facilities, skilled nursing facilities, and home health agencies.

Download the AOTA and APTA Joint Statement and Chartbook at www.aota.org/TOPS and www.apta.org.

Appendix

Functional Scores

The researchers created a standardized functional scale using items from the MDS, IRF-PAI, and OASIS.

Assessment Names

ADL Category	OASIS Item	MDS Item	IRF-PAI Item
Dressing	Dress Upper Body (M1810)	Dressing G0110G1	Dressing – Upper 39D
	Dress Lower Body (M1820)		Dressing – Lower 39E
Walking/ Locomotion	Ambulation/Locomotion (M1860)	Locomotion: On Unit G0110E1 Off Unit G0110F1	Walk/Wheelchair 39L
Toileting	Toileting Hygiene (M1845)	Toilet Use G0110I1	Toileting 39F
Bathing	Bathing (M1830)	Bathing G0120A	Bathing 39C
Transferring	Transferring M1850	Transfer G0110B1	Transfers (Bed, Chair, Wheelchair) 39I
Feeding	Feeding or Eating M1870	Eating G0110H1	Eating 39A



Home Health Deciles

Each decile contains 10% of HH stays with the specified primary diagnosis.

Decile	Total Hours of OT & PT			
	All Diagnoses	Stroke	Congestive Heart Failure	Joint Replacement
1	≤ 1.5	≤ 1.8	≤ 1	≤ 3.8
2	> 1.5 & ≤ 3	> 1.8 & ≤ 3.3	> 1 & ≤ 2.3	> 3.8 & ≤ 4.8
3	> 3 & ≤ 4.3	> 3.3 & ≤ 4.8	> 2.3 & ≤ 3.5	> 4.8 & ≤ 5.5
4	> 4.3 & ≤ 5.3	> 4.8 & ≤ 6	> 3.5 & ≤ 4.5	> 5.5 & ≤ 6.3
5	> 5.3 & ≤ 6	> 6 & ≤ 7.3	> 4.5 & ≤ 5.5	> 6.3 & ≤ 7
6	> 6 & ≤ 7	> 7.3 & ≤ 8.8	> 5.5 & ≤ 6.5	> 7 & ≤ 7.8
7	> 7 & ≤ 8.3	> 8.8 & ≤ 10.5	> 6.5 & ≤ 7.8	> 7.8 & ≤ 8.8
8	> 8.3 & ≤ 10	> 10.5 & ≤ 13	> 7.8 & ≤ 9.5	> 8.8 & ≤ 10
9	> 10 & ≤ 12.5	> 13 & ≤ 16.5	> 9.5 & ≤ 12	> 10 & ≤ 12.3
10	> 12.5 & ≤ 30	> 16.5 & ≤ 29.3	> 12 & ≤ 21.3	> 12.3 & ≤ 30

SNF Deciles

Each decile contains 10% of SNF stays with the specified primary diagnosis.

Decile	Total Hours of OT & PT			
	All Diagnoses	Stroke	Congestive Heart Failure	Joint Replacement
1	<= 6.3	<= 9.3	<= 6.9	<= 3.7
2	> 6.3 & <= 12	> 9.3 & <= 16.8	> 6.9 & <= 12	> 3.7 & <= 7.1
3	> 12 & <= 16.9	> 16.8 & <= 22.8	> 12 & <= 16.8	> 7.1 & <= 10.4
4	> 16.9 & <= 22.3	> 22.8 & <= 29.6	> 16.8 & <= 21.9	> 10.4 & <= 13.7
5	> 22.3 & <= 27.6	> 29.6 & <= 39	> 21.9 & <= 26.5	> 13.7 & <= 17.5
6	> 27.6 & <= 36.4	> 39 & <= 49.9	> 26.5 & <= 34.1	> 17.5 & <= 22.3
7	> 36.4 & <= 46.4	> 49.9 & <= 65.3	> 34.1 & <= 42.4	> 22.3 & <= 27.5
8	> 46.4 & <= 61.8	> 65.3 & <= 88.7	> 42.4 & <= 55.4	> 27.5 & <= 37.9
9	> 61.8 & <= 90.1	> 88.7 & <= 128	> 55.4 & <= 80.1	> 37.9 & <= 58.3
10	> 90.1 & <= 219	> 128 & <= 219.3	> 80.1 & <= 165	> 58.3 & <= 151.9

IRF Deciles

Each decile contains 10% of IRF stays with the specified primary diagnosis.

Decile	Total Hours of OT & PT			
	All Diagnoses	Stroke	Congestive Heart Failure	Joint Replacement
1	<= 12.9	<= 15.1	<= 12.8	<= 12.3
2	> 12.9 & <= 16.5	> 15.1 & <= 19.8	> 12.8 & <= 15.9	> 12.3 & <= 15
3	> 16.5 & <= 19.3	> 19.8 & <= 23.6	> 15.9 & <= 18.1	> 15 & <= 17.3
4	> 19.3 & <= 21.8	> 23.6 & <= 27	> 18.1 & <= 20.5	> 17.3 & <= 19.2
5	> 21.8 & <= 24.4	> 27 & <= 30.6	> 20.5 & <= 22.5	> 19.2 & <= 21.2
6	> 24.4 & <= 27.1	> 30.6 & <= 34.3	> 22.5 & <= 24.5	> 21.2 & <= 23.4
7	> 27.1 & <= 30.2	> 34.3 & <= 39.1	> 24.5 & <= 26.8	> 23.4 & <= 25.9
8	> 30.2 & <= 34	> 39.1 & <= 45.8	> 26.8 & <= 30	> 25.9 & <= 29
9	> 34 & <= 41.1	> 45.8 & <= 54	> 30 & <= 34.7	> 29 & <= 33.3
10	> 41.1 & <= 85	> 54 & <= 85	> 34.7 & <= 55.7	> 33.3 & <= 50.2