Orsini
Specialty Pharmacy Services

Fax To: 877.339.4602 **Phone:** 800.356.5034

EXONDYS 51™ INFUSION ORDER

(Eteplirsen)

1 PATIENT INFORMATION

Patient Name: ____________________

Primary Phone: ____________ DOB: ____________

Allergy: ____________________

Gender: ☐ Male ☐ Female

Patient Weight: ____ Lbs ____ Kg Date Weighed: ________

Diagnosis: *G71.01 Muscular Dystrophy*

2 PRESCRIBER INFORMATION

Prescriber's Name: ____________________

State License #: ____________ NPI #: ____________

Address: ____________________

City, State, Zip: ____________________

Phone: ____________ Fax: ____________

Contact Person: ____________ Phone: ____________

3 CATHETER ACCESS AND FLUSH PROTOCOL

ACCESS TYPE	CATHETER FLUSH ORDERS	PORT OCCLUSION
☐ Peripheral ☐ PORT *(Also include Peripheral IV PRN Port Malfunction)*	**0.9% Saline Flush:** **Dispense:** 30 Days **Refills:** PRN x 1yr Flush line/port with 10mL for patency/SASH protocol. **Heparin Flush:** **Dispense:** 30 Days **Refills:** PRN x 1yr Flush port with ______ mL of Heparin ______ units/mL per SASH protocol.	**CathFlo:** 2mg/2mL as directed. *Pharmacy authorized to dispense upon home health nurse validated port occlusion.* ☐ **Dispense:** 1 Kit **Refills:** PRN x 1yr

4 TREATMENT REGIMEN

Administer pre-medication 30-60 min prior to drug infusion

PRE-MEDICATION

☐ Antipyretic: ____________________ **Qty:** 6 month **Refills:** PRN x 1yr

☐ Antihistamine: ____________________ **Qty:** 6 month **Refills:** PRN x 1yr

☐ EMLA (Lidocaine 2.5%/Prilocaine 2.5%) or ☐ LMX4 cream (Lidocaine 4%) **Qty:** 30 Gram **Refills:** PRN x 1yr ☐ Other: ____________

Apply 1-2 hours before port access.

MEDICATION	ROUTE	DOSE	DIRECTIONS	DAY SUPPLY	REFILLS
Exondys 51™	☒ IV	30 mg/kg	____________mg every week (Supplied as 50mg/mL vials)	30	☒ 1 year
Skilled Nursing Visit	☐ As needed for IV access, administration and proper clinical monitoring		Administration procedures to be followed per pharmacy protocol		

Infusion Volume: Add final drug dose volume to 100-150mL 0.9% saline. ☐ Other: ____________________

Infusion Rate: Over ________ minutes *(Manufacturer recommended infusion duration between 35-60 minutes)*
(Please Specify)

Post Infusion: Flush IV with 25mL 0.9% saline at final rate of drug infusion

Vital Signs: At baseline and completion of post infusion flush. ☐ Other: ____________________

☒ **Supplies**: Provide infusion pump if needed, IV Pole, back-up peripheral IV kit and all necessary infusion supplies

5 PROVIDER SIGNATURE

Product Substitution Permitted Signature ____________ Date of Signature ____________

Dispense as Written Signature ____________ Date of Signature ____________

IMPORTANT NOTICE: This facsimile transmission is intended to be delivered only to the named addressee and may contain material that is confidential, privileged, proprietary or exempt from disclosure under applicable law. If it is received by anyone other than the named addressee, the recipient should immediately notify the sender at the address and telephone number set forth herein and obtain instructions as to disposal of the transmitted material. In no event should such material be read or retained by anyone other than the named addressee, except by the express authority of the sender to the named addressee.

Copyright © 2016 Orsini Healthcare. All Rights Reserved.